DHEA Micronized 10 mg/25 mg

Supports Healthy Androgen and Estrogen Levels*



DHEA Micronized 10 mg and 25 mg are available in 60 tablets

Clinical Applications

- » Supports Healthy Androgen and Estrogen Levels*
- » Hormone Precursor to Androgens*
- » Supports Healthy DHEA Levels*

DHEA Micronized features dehydroepiandrosterone (DHEA), a steroid hormone precursor that is converted by the body into androgen, estrogen, and other hormones. DHEA naturally peaks in early adulthood and declines with age.*

Discussion

Dehydroepiandrosterone (DHEA) is a naturally occurring steroid hormone precursor synthesized from cholesterol primarily by the adrenal glands, and it is produced in small amounts in the ovaries, the testes, and the brain. DHEA production gradually increases in adolescence, typically peaking between the ages of 20 and 30; then it slowly declines approximately 2% each year until about age 70, when it slows to 10% to 20% of the amount encountered in a young adult at peak level.*1-5

The body's DHEA level consists of the combined total of DHEA and its sulfated form, DHEA-sulfate (DHEA-S), the most abundant circulating steroid in humans. They are converted via multiple pathways into hormones such as testosterone and estrogen as needed. Supplemental DHEA is aimed at enhancing the DHEA reservoir so that an increased level is available for conversion to androgens, estrogens, and other hormones involved in a wide variety of physiological functions.*1,3-5

In a randomized, placebo-controlled, crossover trial in men (n = 13) and women (n = 17) aged 40 to 70 years, researchers administered a nightly dose of 50 mg of DHEA to test the hypothesis that a decline in DHEA contributes to the shift from anabolism to catabolism that is associated with aging. Within 2 weeks of DHEA replacement, serum levels of DHEA and DHEA-S were restored to levels typically found in individuals in their second decade of life and were sustained throughout the 3-month study period. The levels of the androgens androstenedione, testosterone, and dihydrotestosterone were doubled in female subjects, with only a slight increase in androstenedione in male subjects. DHEA, at the 50-mg dose level, induced a 10% increase in bioavailable insulin-like growth factor (IGF-1), which the authors suggest could result in an improvement in catabolic processes and physical/psychological well-being over time.*

In a follow-up study led by the same researcher as the study above, a larger dose (100 mg) of DHEA over a 6-month period resulted in the elevation of circulating DHEA and DHEA-S. In the female subjects, a biotransformation to androgen levels similar to their younger counterparts was observed. IGF-1 levels were analyzed in both genders, with the male group showing a higher increase in lean body mass and muscle strength,

suggesting a gender-specific response. Further research is warranted to explore gender and other potential contributing factors.*7

In a randomized trial with healthy subjects (n = 280) aged 60 to 79 over a 1-year period, subjects were given 50 mg of DHEA daily to re-establish DHEA levels. Results indicated that replacement therapy normalized a wide variety of age-related physiological functions and supported wellbeing. The authors suggested a need for further analysis of data.*8

A meta-analysis of 25 randomized, double-blind, placebo-controlled studies in elderly men evaluated the signs and symptoms of low levels of age-related DHEA and the effect of DHEA supplementation. DHEA levels were found to be correlated with many representative age-related phenomena. However, supplementation was found to induce only a small, albeit significant, effect on body composition that was dependent on conversion into the bioactive metabolites of androgen or estrogen.*4

A recent review explored the role of DHEA as an indirect intermediate to androgens and estrogens in relation to varying health conditions in the body and included an overview of the mechanisms of action. The authors concluded that emerging research continues to confirm that DHEA supplementation may be beneficial for older individuals or those with endocrine deficiencies. They suggested that more studies in large populations are needed to validate immunomodulatory and cardiovascular benefits, further assess positive effects on lean muscle and fat mass, and evaluate administration protocols in postmenopausal women.*3

Decades of evidence support DHEA supplementation to restore the DHEA reservoir to peak or near-peak levels in older populations. However, additional well-designed, large-scale trials are needed to clarify guidelines for clinical use.*3.5

DHEA Micronized 10 mg Supplement Facts

Serving Size: 1 Tablet

	Amount Per Serving	%Daily Value
Micronized Dehydroepiandrosterone (DHEA)	10 mg	**
** Daily Value not established.		

Other Ingredients: Dicalcium phosphate dihydrate, microcrystalline cellulose, croscarmellose sodium, ascorbyl palmitate, hydroxypropyl cellulose, silica, and coating (hydroxypropyl methylcellulose and medium-chain triglycerides).

DIRECTIONS: Take one tablet twice daily, or as directed by your healthcare professional. Individuals taking medication should discuss potential interactions with their healthcare professional. Do not use if tamper seal is damaged.

WARNING: NOT FOR USE BY INDIVIDUALS UNDER THE AGE OF 18 YEARS. DO NOT USE IF PREGNANT OR NURSING. Consult a physician or licensed, qualified healthcare professional before using this product if you have, or have a family history of, prostate cancer, prostate enlargement, heart disease, low "good" cholesterol (HDL), or if you are using any other dietary supplement, prescription drug, or over-the-counter drug. Do not exceed recommended serving. Exceeding recommended serving may cause serious adverse health effects. Possible side effects include acne, hair loss, hair growth on the face (in women), aggressiveness, irritability, and increased levels of estrogen. Discontinue use and call a physician or licensed, qualified healthcare professional immediately if you experience rapid heartbeat, dizziness, blurred vision, or other similar symptoms. KEEP OUT OF REACH OF CHILDREN.

FORMULATED TO EXCLUDE: Wheat, gluten, yeast, soy, animal and dairy products, fish, shellfish, peanuts, tree nuts, egg, sesame, ingredients derived from genetically modified organisms (GMOs), artificial colors, and artificial sweeteners.

DHEA Micronized 25 mg Supplement Facts

Serving Size: 1 Tablet

	Amount Per Serving	%Daily Value
Micronized Dehydroepiandrosterone (DHEA)	25 mg	**
** Daily Value not established.		

Other Ingredients: Dicalcium phosphate dihydrate, microcrystalline cellulose, croscarmellose sodium, ascorbyl palmitate, hydroxypropyl cellulose, silica, and coating (hydroxypropyl methylcellulose and medium-chain triglycerides).

DIRECTIONS: Take one tablet twice daily, or as directed by your healthcare professional. Individuals taking medication should discuss potential interactions with their healthcare professional. Do not use if tamper seal is damaged.

WARNING: NOT FOR USE BY INDIVIDUALS UNDER THE AGE OF 18 YEARS. DO NOT USE IF PREGNANT OR NURSING. Consult a physician or licensed, qualified healthcare professional before using this product if you have, or have a family history of, prostate cancer, prostate enlargement, heart disease, low "good" cholesterol (HDL), or if you are using any other dietary supplement, prescription drug, or over-the-counter drug. Do not exceed recommended serving. Exceeding recommended serving may cause serious adverse health effects. Possible side effects include acne, hair loss, hair growth on the face (in women), aggressiveness, irritability, and increased levels of estrogen. Discontinue use and call a physician or licensed, qualified healthcare professional immediately if you experience rapid heartbeat, dizziness, blurred vision, or other similar symptoms. KEEP OUT OF REACH OF CHILDREN.

FORMULATED TO EXCLUDE: Wheat, gluten, yeast, soy, animal and dairy products, fish, shellfish, peanuts, tree nuts, egg, sesame, ingredients derived from genetically modified organisms (GMOs), artificial colors, and artificial sweeteners.

References

- Rutkowski K, Sowa P, Rutkowska-Talipska J, et al. *Drugs*. 2014;74(11):1195-1207. doi:10.1007/s40265-014-0259-8.
- Labrie F, Bélanger A, Cusan L, et al. J Clin Endocrinol Metab. 1997;82(8):2396-2402. doi:10.1210/jcem.82.8.4160.
- Sahu P, Gidwani B, Dhongade HJ. Steroids. 2020;153:108507. doi:10.1016/j.steroids.2019.108507.
- Corona G, Rastrelli G, Giagulli VA, et al. J Clin Endocrinol Metab. 2013;98(9):3615-3626. doi:10.1210/jc.2013-1358.
- Traish AM, Kang HP, Saad F, et al. J Sex Med. 2011;8(11):2960-2982; quiz 2983. doi: 10.1111/j.1743-6109.2011.02523.x.
- Morales AJ, Nolan JJ, Nelson JC, et al. J Clin Endocrinol Metab. 1994;78(6):1360-1367. doi:10.1210/icem.78.6.7515387.
- Morales AJ, Haubrich RH, Hwang JY, et el. Clin Endocrinol (Oxf). 1998;49(4):421-432. doi:10.1046/j.1365-2265.1998.00507.x.
- Baulieu EE, Thomas G, Legrain S, et al. Proc Natl Acad Sci U S A. 2000;97(8):4279-4284. doi:10.1073/pnas.97.8.4279.

